

**MISCELLANEOUS NEW NAMES AND COMBINATIONS IN
SCROPHULARIACEAE - MANULEAE**

O.M. HILLIARD

Manulea derustiana is newly described, and a ditypic genus, *Reyemia*, is described with a new species, *R. chasmanthiflora*, as its type, and *Zaluzianskya nemesioides* is transferred to it.

Manulea derustiana Hilliard, **species nova** a *M. latiloba* Hilliard calycis lobis longioribus et proportione angustioribus (anticis 2.5–3 x 0.8mm, nec 1.5–2 x 0.6–1mm), dorso marginibusque pilis sphaerocephalis dispersis usque ad 0.25mm longis (nec pilis sphaerocephalis sparsis ± sessilibus) praeditis, corollae tubo c.4mm longo (nec 2–2.5mm) differt.

Type: Cape, Oudtshoorn distr., 3322 CB, De Rust, north of Le Roux station, 1800ft, 5 viii 1990, *Viviers & Vlok* 479 (holo. E, iso. K, NBG, S).

Reyemia Hilliard, **genus novum** a *Zaluzianskya* F.W. Schmidt floribus distincte pedicellatis et laxe paniculatis (nec floribus sessilibus vel subsessilibus in racemos terminales dispositis), corollae limbo resupinato, pilis clavatis in postico faucis latere (nec fauce glabro etsi ore saepe pilis circumcincto), staminibus 2 et staminodiis 2 (nec staminibus 4 vel 2, staminodiis absentibus) differt. A *Phyllopodio* Benth. forma corollae et limbo corollae resupinato (labio antico integro, labio postico 4-lobato, nec lobo antico 3-lobed, limbo postico bilobato, staminibus 2 posticis fertilibus et staminodiis 2 (nec staminibus plerumque 4 fertilibus sed specie unica *Phyllopodii* staminibus posticis abortis), filamentis ad basin corollae tubi decurrentibus (nec breviter decurrentibus tantum) differt.

Type species: *Reyemia chasmanthiflora* Hilliard.

The generic name commemorates Dr H Meyer, who practised medicine in Calvinia in the 1860s and whose collections from the Hantam Mountains are the basis of many names, including *R. nemesioides*.

Reyemia chasmanthiflora Hilliard, **species nova** a *R. nemesioide* (Diels) Hilliard corollae tubo multo longiore (c.27–30mm longo, nec 7–8.5mm) et lobis angustioribus (lobo antico 7.5 x 1.5mm, nec 3.5–6 x 3.6 x 5mm, lobis posticis c.7 x 2.2mm, nec 3.5 x 3mm) facile distinguitur.

Type: Cape, Calvinia div., 3120 CB, farm Annexe Kransfontein 721, 1123m, 2 ix 1986, *Cloete & Haselau* 172 (holo. STE).

Reyemia nemesioides (Diels) Hilliard, **comb. nov.**

Type: Cape, Calvinia div., Hantam Mountains, *Meyer* s.n. (holo. B+).

Syn.: *Zaluzianskya nemesioides* Diels in Bot. Jahrb. 23: 482 (1896).