

- A. Ophiopogon** Lévl. in Fedde, Rep. Spec. Nov. xii. 184 (Mai. 1913);
ex Takeda in Jap. Bot. Mag. Tok. xxvii. 215 (Nov. 1913)—
Korea (Ins. Quelpaert).—Sp. propr. (ut videtur).
A. praelatitium Lévl. in Fedde, Rep. Spec. Nov. xii. 288 (1913)—Yun-
nan - - - - - = **Bulleyanum** Diels
A. Simethis Lévl. et Giraud. in Fedde, Rep. Spec. Nov. xii. 288 (1913)
—Yunnan - - - - - = **macranthum** Baker
A. Taqueti Lévl. in Fedde, Rep. Spec. Nov. v. 283 (1908)—Korea
(Ins. Quelpaert) - - - - - = **japonicum** Thunb.
A. tchongchanense Lévl. l.c. vi. 263 (1909).—*A. odorum* Lévl. Cat. Pl.
Yunn. 163 (1915-17), non L.—Yunnan - - - - -
= **Bulleyanum** var. *tchongchanense* A.-S.

New Species of Viola from South-East Asia.

BY

WILHELM BECKER.

Viola Unwinii W. Bckr., sp. nov.

Sect. *Nomimium* Ging.; ex affinitate *V. annamensis* Baker f., quae foliis ovato-oblongis basi plus minusve cuneatis margine plus minusve serratis differt.

Planta acaulis, glaberrima, 8 cm. alta, stolonifera. Rhizoma verticale, flexuosum, non crassum, articulatum, radicellatum. Stolones 1 (vel plures?) tenues, plus minusve ascendentes, ad finem foliati et denique radicantes rosulam formantes. Stipulae lanceolatae, 1-1.5 cm. longae, virides, plus minusve dense longifimbriatae; fimbriae usque ad 2 mm. longae. Folia basi plane cordata, ovato-lanceolata, 2.5-4.5 cm. longa et 1.2-2.2 cm. lata, acutiuscula, plane crenato-serrata (circ. 10-12 crenis), glaberrima, longe petiolata. Flores albi, folia non superantes; pedicelli in parte superiore bracteolis circ. 8 mm. longis remote longifimbriatis muniti; sepala anguste lanceolata, 6 mm. longa, brevissime appendiculata; petala apice rotundata, superiora obovato-oblonga 7 mm. longa et 3.75 mm. lata, lateralia oblonga 8 mm. longa et 3 mm. lata ebarbata; pet. infimum abbreviatum, 7.5 mm. longum (calc. inclus.), naviculare, angustatum, acutiusculum, breviter calcaratum; calcar crassum, 2 mm. longum et latum; ovarium conoideum; stylus basi geniculatus, apice horizontaliter deplanatus, submarginatus, brevissime rostellatus.

Burma: Mt. Victoria, Kanpetlet, 6250 ft., on steep mossy banks; coll. R. Unwin 14.4.1926, 3017. Typus in Herb. Royal Bot. Gard. Edinb., Cotyp. in W. Bckr. Herb. Viol.

Viola nuda W. Bckr., sp. nov.

Sect. *Nomimium* Ging., ex affinitate *V. Thomsonii* Oudem.; foliis glaberrimis distincte late ovatis non acuminatis glaberrimis, sepalis latioribus et stipulis plus minusve viridibus distincte diversa.

Planta acaulis, glaberrima, stolonifera, 7–13 cm. alta. Rhizoma verticale, circ. 2 mm. crassum, subflexuosum, distincte articulatum, longe radicellatum. Stolones 1 (vel pauci), ascendentes, adversus finem foliati, verosimiliter posterius radicales, 10–15 cm. longi. Stipulae non adnatae, lanceolatae, virides, circ. 1.5 cm. longae, subdense longifimbriatae. Folia persistentia longe petiolata, circ. 5–6 cm. longa et 3–4 cm. lata, basi profunde et subanguste cordata, ceterum distincte ovata, subacutiuscula, glaberrima, plane crenato-serrata (circ. 20 crenis), remote 5-nervia; folia novella eadem forma, minora. Flores pauci, folia non superantes, albi, ad basin rubro-purpureo-lineati et maculati; pedicelli bracteolis 8 mm. longis, in parte superiore insertis instructi; sepala ovata, 5 mm. longa et 2–2.5 mm. lata, etiam latiora et plus minusve crenulata, breviter appendiculata; petala superiora obovata, lateralia obovato-oblonga subbarbulata, 1 cm. longa; pet. infimum abbreviatum circ. 8 mm. longum naviculare, acutiusculum; stylus basi geniculatus, apice horizontaliter deplanatus, submarginatus, brevissime rostellatus.

W. Yunnan: Shweli-Salween divide, 8000 ft. (Lat. 25° 20' N., Long. 98° 58' E.), on banks and slopes in thickets; coll. George Forrest IV. 1924, 24019. Typus in Herb. Royal Bot. Gard. Edinb.