

GEOGRAPHICAL DISTRIBUTION OF MICROCYCLIC SPECIES OF THE THREE RUST GENERA, PUCCINIA, UROMYCES AND PHRAGMIDIUM IN HOKKAIDO, JAPAN*

N. HIRATSUKA**

ABSTRACT. Geographical data is presented for the microcyclic species of the rust genera, *Puccinia*, *Uromyces* and *Phragmidium* (Uredinales), that occur in Hokkaido, Japan. The distribution of these species in other parts of the Japanese Archipelago is tabulated, and the species are classified into six phytogeographical elements.

It is an interesting matter to compare the rust-flora in Hokkaido, northern district of Japan, with that of other districts in the world, and to consider their relationships. In this paper data on the geographical distribution of microcyclic species of the three rust genera, *Puccinia*, *Uromyces* and *Phragmidium* in Hokkaido, Japan is presented.

The following enumeration is a general list of the microcyclic species of the three genera of Uredinales in Hokkaido, northern district of Japan, based upon the data in a recent paper on the Uredinales of Hokkaido (Hiratsuka *et al.*, 1986).

The specimens of these rust fungi are now mostly preserved in the Hiratsuka Herbarium (Tokyo).

1. *Puccinia adenophorae* Dietel on *Adenophora remotiflora* & *A. triphylla* var. *japonica*.
2. *Puccinia adoxae* Hedwig f. on *Adoxa moschatellina*.
3. *Puccinia alpina* Fuckel on *Viola vaginata*.
4. *Puccinia anemones-raddeanae* S. Ito on *Anemone raddeana*.
5. *Puccinia arenariae* (Schum.) Winter on *Moehringia lateriflora*, *Spergula arvensis*, *Stellaria diversiflora* & *S. media*.
6. *Puccinia artemisiae-keiskeanae* Miura on *Artemisia keiskeana*.
7. *Puccinia asarina* Kunze on *Asarum heterotropoides*.
8. *Puccinia austroberingiana* (Savile) Savile on *Saxifraga fusca* var. *divaricata*.
9. *Puccinia campanulae* Carmich. in Berk. on *Campanula lasiocarpa*.
10. *Puccinia chrysosplenii* Grev. on *Chrysosplenium flagelliferum*.
11. *Puccinia circaeae* Pers. on *Circaea caulescens* var. *glabra* & *C. erubescens*.
12. *Puccinia cnici-oleracei* Pers. on *Achillea ptarmica* var. *macrocephala*, *Artemisia japonica*, *A. japonica* var. *macrocephala*, *A. montana*, *A. princeps*, *Cacalia hastata* subsp. *orientalis*, *Chrysanthemum parthenium*, *Ch. praealtum*, *Matricaria chamomilla* & *Sonchus brachyotus*.
13. *Puccinia conglomerata* (Strauss) Röhling on *Petasites japonicus* var. *giganteus*.
14. *Puccinia epilobii* DC. on *Epilobium hornemanni* var. *foucaudianum*.
15. *Puccinia exhausta* Dietel on *Clematis stans*.
16. *Puccinia fergussoni* Berk. & Broome on *Viola hirtipes* & *V. selkirkii*.
17. *Puccinia fusispora* P. & H. Sydow on *Urtica platyphylla*.
18. *Puccinia glechomatis* DC. on *Glechoma hederacea* var. *grandis*.
19. *Puccinia glomerata* Grev. on *Cacalia auriculata* var. *kamtschatica* & *Ligularia hodgsoni*.
20. *Puccinia hakkodensis* Hirats. f. on *Clintonia udensis*.

*Contributions from the Tottori Mycological Institute, No 211.

**Tottori Mycological Institute, Kokoge-221, Tottori, 689-11, Japan.

21. *Puccinia haleniae* Arthur & Holway on *Halenia corniculata* & *Swertia tetrapetala*.
- 22a. *Puccinia heucherae* (Schw.) Dietel var. *japonica* Savile on *Saxifraga japonica*.
- 22b. *Puccinia heucherae* var. *minor* Savile on *Mitella nuda*.
23. *Puccinia horiana* Henn. on *Chrysanthemum morifolium* var. *sinense*.
24. *Puccinia lantanae* Farlow on *Justicia procumbens* var. *leucantha*.
25. *Puccinia majanthemi* Dietel on *Convallaria keiskei*, *Majanthemum bifolium*, *M. dilatatum*, *Polygonatum maximowiczii*, *Trillium kamtschaticum* & *T. tschonoskii*.
26. *Puccinia melanoplaca* P. & H. Sydow on *Patrinia gibbosa*.
27. *Puccinia ornata* Arthur & Holway on *Rumex obtusifolius*.
28. *Puccinia parkerae* Dietel & Holway on *Ribes japonicum* & *R. sachalinense*.
29. *Puccinia patriniae* Henn. on *Patrinia villosa*.
30. *Puccinia polemonii* Dietel & Holway on *Polemonium acutiflorum* var. *laxiflorum* & *P. yezoense*.
31. *Puccinia porphyrogenita* Curtis on *Chamaepericlimenum canadense*.
32. *Puccinia ribis* DC. on *Ribes latifolium* & *R. triste*.
33. *Puccinia saxifragae-micranthae* Barclay on *Saxifraga sachalinensis*.
34. *Puccinia schizonepetae* Tranzschel on *Nepeta subsessilis* var. *yezoensis*.
35. *Puccinia singularis* Magnus on *Anemone altaica*.
36. *Puccinia tiarellaecola* Hirats. f. on *Tiarella polyphylla*.
37. *Puccinia togashiana* Hirats. f. on *Thalictrum aquilegifolium* & *Th. thunbergii* var. *majus*.
38. *Puccinia tokubuchii* S. Ito on *Patrinia gibbosa*.
39. *Puccinia trollii* Karsten on *Trollius riederianus* var. *japonicus*.
40. *Puccinia umbilici* Guepin on *Rhodiola rosea* var. *tashiroi*.
41. *Puccinia ustalis* Berk. on *Trautvetteria japonica*.
42. *Puccinia veronicae* Schröter on *Veronica nipponica*.
43. *Puccinia veronicarum* DC. on *Veronicastrum sibiricum* var. *japonicum*.
44. *Puccinia vesiculosa* Schlechtendal on *Anemone narcissiflora*.
45. *Puccinia virgae-aureae* (DC.) Lib. on *Solidago virgae-aurea* & *S. virgae-aurea* subsp. *leiocarpa*.
46. *Puccinia waldsteiniae* Curtis on *Waldsteinia ternata*.
47. *Uromyces cacaliae* (DC.) Unger on *Cacalia auriculata* var. *bulbifera* & *C. auriculata* var. *kamtschatica*.
48. *Uromyces flectens* Lagerh. on *Trifolium repens*.
49. *Uromyces gageae* G. Beck on *Gagea lutea* & *Lloydia triflora*.
50. *Uromyces miurae* P. & H. Sydow on *Fritillaria kamtschaticensis*.
51. *Uromyces peracarpae* S. Ito & Tochinai on *Percarpa circaeoides*.
52. *Uromyces rudbeckiae* Arthur & Holway on *Solidago virga-aurea* subsp. *asiatica*.
53. *Uromyces sommerfeltii* Hylander, Jørstad & Nannfeldt on *Solidago virga-aurea* subsp. *leiocarpa*.
54. *Phragmidium alpinum* Hirats. f. on *Rubus pedatus*.
55. *Phragmidium kamtschatkae* (F. W. Anderson) Arthur & Cummins on *Rosa rugosa*.
56. *Phragmidium miyabeana* S. Ito & Hirats. f. on *Sieversia pentapetala*.

DISTRIBUTION OF SPECIES IN THE JAPANESE ARCHIPELAGO

Table 1 presents the distribution of the microcyclic species of *Puccinia*, *Uromyces* and *Phragmidium* recorded from Hokkaido in relation to the other districts in the Japanese Archipelago.

PHYTOGEOGRAPHICAL CLASSIFICATION

The 56 species (including two varieties) of microcyclic *Puccinia*, *Uromyces* and *Phragmidium* found in Hokkaido may be classified in the following six groups.

TABLE I
Japanese Archipelago
Japan

	Sakhalin	Hokkaido	Northern Honshu	Central Honshu	Western Honshu	Shikoku	Kyushu	Ryukyu Islands	Formosa (Taiwan)
<i>Puccinia adenophorae</i>	+	+	+	+ ^a	+ ^a		+ ^a		+ ^a
<i>P. adoxae</i>		+							
<i>P. alpina</i>		+	+	+	+				
<i>P. anemones-raddeanae</i>	+	+							
<i>P. arenariae</i>	+	+	+	+	+	+	+		+ ^a
<i>P. artemisiae-keiskeanae</i>		+	+	+	+	+			
<i>P. asarina</i>	+	+	+	+					
<i>P. austroberingiana</i>	+	+	+	+	+		+		
<i>P. campanulae</i>		+ ^m	+ ^m						
<i>P. chrysosplenii</i>	+	+		+	+	+	+		
<i>P. circaeae</i>	+	+	+	+					
<i>P. cnici-oleracei</i>	+	+	+	+	+	+	+	+	+
<i>P. conglomerata</i>	+	+	+						
<i>P. epilobii</i>		+ ^m							
<i>P. exhausta</i>		+	+	+	+			+	+
<i>P. fergussoni</i>	+	+	+						
<i>P. fusispora</i>		+							
<i>P. glechomatis</i>		+	+	+	+	+	+	+	+
<i>P. glomerata</i>	+	+							
<i>P. hakkodensis</i>	+	+	+	+					
<i>P. haleniae</i>	+	+		+ ^a					
<i>P. heucherae</i> var. <i>japonica</i>		+	+ ^m						
<i>P. heucherae</i> var. <i>minor</i>		+							
<i>P. horiana</i>		+	+	+	+	+	+	+	
<i>P. lantanae</i>		+	+	+	+	+	+	+	+
<i>P. majanthemi</i>	+	+	+ ^m	+ ^m					
<i>P. melanoplaca</i>		+	+	+					
<i>P. ornata</i>		+							
<i>P. parkerae</i>	+	+	+	+		+ ^m			
<i>P. patriniae</i>		+	+	+	+	+	+	+	+
<i>P. polemonii</i>	+	+ ^m		+ ^a					
<i>P. porphyrogenita</i>	+	+ ^m	+ ^m	+ ^a					
<i>P. ribis</i>	+	+							
<i>P. saxifragae-micranthae</i>		+							
<i>P. schizonepetae</i>		+		+ ^a		+ ^m			
<i>P. singularis</i>		+							
<i>P. tiarellaecola</i>	+	+	+	+					
<i>P. togashiana</i>		+	+	+					
<i>P. tokubuchii</i>		+							
<i>P. trollii</i>		+ ^m							
<i>P. umbilici</i>		+	+ ^m	+ ^m					+ ^a
<i>P. ustalis</i>	+	+	+ ^m	+ ^m					+ ^a
<i>P. veronicae</i>		+	+ ^m						
<i>P. veronicarum</i>		+	+	+ ^m					
<i>P. vesiculosa</i>	+	+ ^m	+ ^a	+ ^a					
<i>P. virgae-aureae</i>	+	+ ^m	+ ^a	+ ^a					
<i>P. waldsteiniae</i>	+	+		+					
<i>Uromyces calaciae</i>	+	+							
<i>U. flectens</i>		+	+	+					
<i>U. gageae</i>		+							
<i>U. miurae</i>	+	+	+	+ ^m					
<i>U. peracarpae</i>	+	+							
<i>U. rudbeckiae</i>	+	+	+	+	+	+	+		+
<i>U. sommerfeltii</i>	+	+ ^m	+ ^m	+ ^m	+ ^m				
<i>Phragmidium alpinum</i>	+ ^m	+ ^m		+ ^m					
<i>Ph. kamtschatkae</i>	+	+							
<i>Ph. miyabeana</i>	+ ^m	+ ^m	+ ^m	+ ^m					

+^m denotes records confined to alpine or subalpine regions of high mountains.

1. CIRCUMPOLAR ELEMENTS

<i>Puccinia adoxae</i>	<i>P. fergussoni</i>
<i>P. arenariae</i>	<i>P. ribis</i>
<i>P. asarina</i>	<i>P. umbilici</i>
<i>P. campanulae</i>	<i>P. ustalis</i>
<i>P. circaeae</i>	<i>P. vesiculosa</i>
<i>P. conglomerata</i>	<i>P. virgae-aureae</i>
<i>P. epilobii</i>	<i>Uromyces sommerfeltii</i>

2. EUROPEAN AND EASTERN ASIATIC ELEMENTS

<i>Puccinia alpina</i>	<i>P. veronicae</i>
<i>P. chrysosplenii</i>	<i>P. veronicarum</i>
<i>P. glechomatis</i>	<i>Uromyces cacaliae</i>
<i>P. glomerata</i>	<i>U. gageae</i>
<i>P. singularis</i>	<i>Phragmidium kamtschatkae</i>
<i>P. trollii</i>	

3. ELEMENTS OF HIMALAYA, CONTINENTAL CHINA AND JAPAN

Puccinia saxifragae-micranthae

4. NORTH AMERICA AND EASTERN ASIATIC ELEMENTS

<i>Puccinia austroberingiana</i>	<i>P. polemonii</i>
<i>P. haleniae</i>	<i>P. porphyrogenita</i>
<i>P. heucherae</i> var. <i>minor</i>	<i>P. waldsteiniae</i>
<i>P. ornata</i>	<i>Uromyces miurae</i>
<i>P. parkerae</i>	<i>U. rudbeckiae</i>

5. EASTERN ASIATIC ELEMENTS

a. Elements of Maritime Provinces of USSR and Japan

<i>Puccinia anemones-raddeanae</i>	<i>P. togashiana</i>
<i>P. hakkodensis</i>	<i>Uromyces peracarpae</i>
<i>P. majanthemi</i>	<i>Phragmidium alpinum</i>
<i>P. tiarellaecola</i>	<i>Ph. miyabeaenum</i>

b. Elements of Continental China, Maritime Provinces of USSR and Japan

<i>Puccinia fusispora</i>	<i>P. schizonepetae</i>
---------------------------	-------------------------

c. Elements of Continental China, Taiwan (China), Korea and Japan

<i>Puccinia adenophorae</i>	<i>P. patriniae</i>
<i>P. artemisiae-keiskeanae</i>	

d. Elements of Java, Philippines, Continental China, Korea, Maritime Provinces of USSR and Japan

Puccinia exhausta

e. Endemic elements:

i. Species peculiar to Hokkaido

Puccinia tokubuchii

ii. Species peculiar to Hokkaido and Honshu

Puccinia heucherae var. *japonica* *P. melanoplaca*

6. COSMOPOLITAN ELEMENTS

<i>Puccinia cnici-oleracei</i>	<i>P. lantanae</i>
<i>P. horiana</i>	<i>Uromyces flectens</i>

REFERENCES

- HIRATSUKA, N. (1935). A contribution to the knowledge of the rust-flora in the alpine regions of high mountains in Japan. *Mem. Tottori Agr. Coll.* 3:125-247.
- (1936). Ueber einige interessante oder für neue Rostpilze. *Ann. Mycol.* 27:278-280.
- (1960). A provisional list of Uredinales of Japan Proper and the Ryukyu Islands. *Sci. Bull. Agr. Div. Univ. Ryukyus.* 7:189-314.
- (1973). Revision of taxonomy of the genus *Uromyces* in Japanese Archipelago. *Rep. Tottori Mycol. Inst.* 10:1-98.
- (1976). Microcyclic species of *Puccinia* in the Japanese Archipelago. *Ibid.* 14:1-77.
- (1980). A taxonomic revision of the autoecious species of *Puccinia* parasitic on the Compositae in the Japanese Archipelago. *Ibid.* 18:1-52.
- , HIRATSUKA, T. & HIRATSUKA, K. (1986). Uredinales of Hokkaido. *Ibid.* 24:203-233.
- & HIRATSUKA, Y. (1958). Uredinales found in the Daisetsu National Park, Hokkaido, Japan. *Sydowia* 12:363-385.
- , KANEKO, S. & NISHIGAKI, H. (1980). A taxonomic revision of the species of *Phragmidium* in the Japanese Archipelago. *Rep. Tottori Mycol. Inst.* 18:53-88.
- & HASEBE, S. (1978). A taxonomic revision of the species of *Puccinia* parasitic on the Liliales (Liliaceae, Amaryllidaceae, Dioscoreaceae and Iridaceae) in the Japanese Archipelago. *Ibid.* 16:1-36.